



# MOVICOL<sup>®</sup> Junior

## Lemon-Lime Flavour

Please read this leaflet carefully before taking your medicine.

If you have any questions or are not sure about anything, ask your doctor or pharmacist.

### What is this medicine?

The name of this medicine is Movicol Junior Lemon-Lime.

Active ingredients per sachet/per 62.5 mL after preparation			
Macrogol 3350	6.563 g	Sodium bicarbonate	89.3 mg
Sodium chloride	175.4 mg	Potassium chloride	23.3 mg
Contains 93.4 mg of sodium and 13 mg of potassium			

Movicol Junior Lemon-Lime also contains lemon-lime flavour, and acesulfame potassium as a sweetener.

Movicol Junior Lemon-Lime is lactose-free, gluten-free and sucrose-free.

Each sachet contains 6.9 grams of Movicol Junior Lemon-Lime powder.

You add water to the powder to make a drink.

### What is Movicol Junior Lemon-Lime used for?

Movicol Junior Lemon-Lime is used in children aged 2 years and older, for the treatment of chronic constipation. It can also be used in adults who need a smaller dose. Constipation is the "less frequent than normal" passing of firm hard stools. Movicol Junior Lemon-Lime helps you to have a comfortable bowel movement even if you have been constipated for a long time.

#### Chronic Constipation in Children:

Most normal children will occasionally experience constipation, which will normally require no more than a healthy diet, plenty of exercise, regular toilet use and, sometimes, occasional use of laxatives. However, a small proportion of children will pass stools less frequently than 3 times per week, with excessive straining and discomfort or pain at these times. If your child has this more severe type of constipation your doctor needs to be involved in making a supervised plan of treatment for your child over the next 6–12 months. Treatments may require daily use of a product such as Movicol Junior Lemon-Lime which can keep stools of normal consistency and restore a normal pattern of passing stools. Movicol Junior Lemon-Lime is also used in children aged 2 years and older for the treatment of faecal impaction (as determined by your doctor), and for the prevention of recurrence of faecal impaction.

#### Before you take Movicol Junior Lemon-Lime

Do not take Movicol Junior Lemon-Lime if your doctor has told you that you have:

- an obstruction in your intestine (gut)
- a perforated gut wall
- severe inflammatory bowel disease, like ulcerative colitis, Crohn's disease, or toxic megacolon
- paralysis of the bowel.
- an allergy to macrogol or any of the ingredients.

#### Pregnancy and breast-feeding

If you are pregnant talk to your doctor before you take Movicol Junior Lemon-Lime.

Movicol Junior Lemon-Lime can be taken while breast-feeding.

#### Other Medications

Laxative products such as Movicol Junior Lemon-Lime have the potential to interact with other medications, by altering their absorption. Close monitoring of the effects of your medications may be necessary when you commence or cease to take Movicol Junior Lemon-Lime regularly. It is important that you should discuss this with your doctor.

If you need to thicken fluids in order to swallow them safely, Movicol Junior Lemon-Lime may counteract the effect of the thickener.

#### How to take Movicol Junior Lemon-Lime

##### Chronic Constipation/Prevention of recurrence of faecal impaction:

**Children aged 2-5 years:** The usual starting dose is 1 sachet daily.

**Children aged 6-11 years:** The usual starting dose is 2 sachets daily. The dose should be adjusted up or down as required to produce regular soft stools. The maximum dose needed does not normally exceed 4 sachets daily.

Use in children aged 2 years and older should be limited to 12 weeks except under medical supervision. Movicol Junior Lemon-Lime is not recommended for children below 2 years of age.

**Adults and children 12 years and older:** The recommended dose of Movicol Junior Lemon-Lime is 2 sachets a day. This can be increased up to 6 sachets daily if necessary, depending on the severity of your constipation. For chronic constipation the dose may be reduced to 1 sachet daily, according to individual response. For patients 12 years and older using 2 sachets daily or more, it is recommended to use Movicol Lemon-Lime (full strength).

#### **FAECAL IMPACTION:**

##### **Children (2 – 11 years):**

A course of treatment with Movicol Junior Lemon-Lime is for up to 7 days as follows:

	<b>Number of Movicol Junior Lemon-Lime Sachets</b>						
<b>Age (years)</b>	<b>Day 1</b>	<b>Day 2</b>	<b>Day 3</b>	<b>Day 4</b>	<b>Day 5</b>	<b>Day 6</b>	<b>Day 7</b>
<b>2 - 5</b>	<b>2</b>	<b>4</b>	<b>4</b>	<b>6</b>	<b>6</b>	<b>8</b>	<b>8</b>
<b>6 - 11</b>	<b>4</b>	<b>6</b>	<b>8</b>	<b>10</b>	<b>12</b>	<b>12</b>	<b>12</b>

Treatment can be stopped when the medicine has worked. This is shown by the patient passing large volumes of stool or watery diarrhoea.

It is not recommended to use Movicol Junior Lemon-Lime for the treatment of faecal impaction in children with a heart or kidney condition.

If you take too much Movicol Junior Lemon-Lime and get bad diarrhoea, stop taking Movicol Junior Lemon-Lime until the diarrhoea stops. If constipation recurs, you should check with your doctor or pharmacist. If you are worried, contact your doctor or pharmacist.

#### **General recommendations when taking laxatives**

Drink plenty of water. Increase fibre in diet except in cases of medication-induced constipation. Prolonged use of laxatives is undesirable and may lead to dependence. If symptoms persist seek medical advice.

#### **How to prepare Movicol Junior Lemon-Lime**

Open the sachet and pour contents into a glass. Add 62.5 mL (about 1/4 cup) of water. Stir well until all the powder has dissolved and the Movicol Junior Lemon-Lime solution is clear or slightly hazy, then drink it. If you like, you can add a flavour such as orange cordial to the drink.

For use in faecal impaction, the correct number of sachets can be made up in advance and kept covered and refrigerated for 24 hours. For example 12 sachets can be made up into 750 mL of water and 16 sachets into one litre of water.

#### **What about side effects?**

Like all medicines, Movicol Junior Lemon-Lime can have side effects. Sometimes people have indigestion, stomach ache or rumbles. You may also feel bloated, suffer from wind, feel sick or vomit, may also experience soreness of the anus (bottom) and may have mild diarrhoea when starting to take Movicol Junior Lemon-Lime. Diarrhoea generally get better if you reduce the amount Movicol Junior Lemon-Lime you take. If necessary, discuss reducing the dose with your doctor or pharmacist. Other side effects include allergic reactions which may cause a skin rash, itching, reddening of the skin or a nettle rash. Tell your doctor or pharmacist if you think Movicol Junior Lemon-Lime Junior is causing you any problem.

Tell your doctor immediately and stop taking Movicol Junior Lemon-Lime if you:

- Get a serious allergic reaction which causes difficulty in breathing, or swelling of the face, lips, tongue or throat.
- If you feel weak, breathless, very thirsty with a headache, or get puffy ankles.

#### **How to store Movicol Junior Lemon-Lime**

##### **Storing Movicol Junior Lemon-Lime sachets**

Store the sachets below 25°C. Do not use Movicol Junior Lemon-Lime after the expiry date on the pack.

##### **Storing prepared solution of Movicol Junior Lemon-Lime**

If you do not drink it straightaway, the prepared Movicol Junior Lemon-Lime solution can be kept covered in the fridge (2 - 8°C) for up to 24 hours. After that time, any remaining solution should be discarded.

Keep out of reach of children.

AUST R 109758

Date: October 2020

**Norgine Pty Limited**, Suite 3.01 Building A, 20 Rodborough Rd, Frenchs Forest NSW 2086  
www.movicol.com.au Ph: 1800 766 936 (AUS) Ph: 0800 404 178 (NZ)



MOVICOL, NORGINE and the sail logo are registered trademarks of the Norgine group of companies.